

Brain Connectivity-Informed Adaptive Regularization for Generalized Outcomes

Jaroslav Harezlak, Ph.D.

Professor and Interim Co-Chair
Department of Epidemiology and Biostatistics
Indiana University School of Public Health
Bloomington, IN, USA

May 22, 2020



Outline

- 1 Motivating application
- 2 Brain structure and connectivity
- 3 Regularization methods
- 4 riPEER - ridgified Partially Empirical Eigenvectors for Regression
- 5 Simulation study
- 6 Brain structure and HIV infection
- 7 Discussion

HIV infection study - WUSM

- ① N = 299 HIV-infected individuals: 228 males, 71 females
 - ▶ Duration of infection range: 0 - 33y (mean: 10.2, sd: 8.7)
 - ▶ Age range: 18 - 84 y.o. (mean: 42.3, sd: 16)
- ② Imaging modalities
 - ▶ T1 - anatomy
 - ▶ DTI - structural connectivity

Anatomy and connectivity


1 Anatomy

- ▶ MPRAGE protocol
- ▶ Processing using FreeSurfer software (version 5.1)
- ▶ Desikan-Killiany atlas - 66 cortical regions

2 Structural connectivity


- ▶ DTI and maximal diffusion coherence model
- ▶ Density of connections between each pair of regions

MRI data




MRI-derived data: cortical thickness

- 1 Parcellation of the cortex into 66 regions
- 2 Average cortical thickness



(a) Parcellation of the brain




(b) Cortical thickness


Connections in the brain



Connectivity matrices




(a) Connectivity matrix: subject 1



(b) Connectivity matrix: subject 2

Population connectivity matrix



Questions



① Scientific

- ▶ Are changes in the brain structure associated with the HIV infection?
- ▶ Is there any additional information provided by the structural connectivity?

② Statistical

- ▶ How to deal with the highly correlated predictors in the regression models?
- ▶ How to incorporate the structural connectivity information in the regression models?


Statistical model



$$y = Zb + X\beta + \varepsilon$$

- ❶ y – n -dimensional response (e.g. NP domain score)
- ❷ $Z \in \mathbb{R}^{n \times 66}$ and $X \in \mathbb{R}^{n \times m}$
- ❸ $\varepsilon \sim \mathcal{N}(0, \sigma^2 I_n)$ for some unknown $\sigma^2 > 0$

Statistical model



- 1 y – n -dimensional response (e.g. NP domain score)
- 2 $Z \in \mathbb{R}^{n \times 66}$ and $X \in \mathbb{R}^{n \times m}$
- 3 $\varepsilon \sim \mathcal{N}(0, \sigma^2 I_n)$ for some unknown $\sigma^2 > 0$

Penalized estimation

To find the estimates of b and β , we consider the optimization problem of the form

$$\operatorname{argmin}_{b, \beta} \left\{ \underbrace{\|y - Zb - X\beta\|_2^2}_{\text{model fit term}} + \lambda \underbrace{g(b)}_{\text{penalty on } b} \right\}.$$

- ❶ $g(b) = \sum_i b_i^2 \quad \longrightarrow \text{Ridge estimate}$
- ❷ $g(b) = \sum_i |b_i| \quad \longrightarrow \text{LASSO estimate}$
- ❸ $g(b) = \|Lb\|_2^2 \quad \longrightarrow \text{Generalized ridge}$

T. W. Randolph, J. Harezlak, Z. Feng, Structured penalties for functional linear models – partially empirical eigenvectors for regression, Electronic Journal of Statistics (2012)

Desired property of the estimate, $\hat{b}$

“Stronger connections between the brain regions i and j result in more similar coefficients $\hat{b}_i$ and $\hat{b}_j$.”

Penalty selection

The natural choice of the penalty is

$$g(b) = \sum_{i,j} a_{ij} (b_i - b_j)^2.$$

- ❶ $d_i := \sum_k A_{ik}.$
- ❷ $D := \text{diag}(d_1, \dots, d_{66})$
- ❸ $Q := D - A$ [Laplacian of A]

Then:

$$\sum_{i,j} a_{ij} (b_i - b_j)^2 = b^T Q b.$$

Connections with the linear mixed models (LMM)

Our objective function becomes

$$\operatorname{argmin}_{b, \beta} \left\{ \|y - Zb - X\beta\|_2^2 + \lambda b^T Q b \right\}.$$

This optimization problem is “equivalent” to the LMM formulation

- ❶ $y = Zb + X\beta + \varepsilon$, where β is a vector of fixed effects and b a vector of random effects,
- ❷ $\varepsilon \sim \mathcal{N}(0, \sigma^2 I)$,
- ❸ $b \sim \mathcal{N}(0, \sigma_b^2 Q^{-1})$,
- ❹ λ, σ and σ_b are connected via $\lambda = \sigma^2 / \sigma_b^2$.

Selection of the regularization parameter



IDEA: define $\hat{\lambda}$ as $\hat{\lambda} = \frac{\hat{\sigma}^2}{\hat{\sigma}_b^2}$ where $\hat{\sigma}^2$ and $\hat{\sigma}_b^2$ are maximum likelihood estimates from the corresponding linear mixed model



PROBLEM: Laplacian is not an invertible matrix, which is required in computation

The method

riPEER (ridgified Partially Empirical Eigenvectors for Regression)

$$\begin{bmatrix} \hat{b}^{\text{rP}} \\ \hat{\beta}^{\text{rP}} \end{bmatrix} := \underset{b, \beta}{\operatorname{argmin}} \left\{ \|y - Zb - X\beta\|_2^2 + \underbrace{\lambda_Q b^T Q b}_{\text{graph part}} + \underbrace{\lambda_R \|b\|_2^2}_{\text{ridge part}} \right\}$$




Figure 3: Different shapes of the set $\{b : \lambda_Q b^T Q b + \lambda_R \|b\|_2^2 \leq 1\}$ for $p = 2$.

Connections with the linear mixed models (LMM)

riPEER (ridgified Partially Empirical Eigenvectors for Regression)


$$\begin{bmatrix} \hat{b}^{\text{rP}} \\ \hat{\beta}^{\text{rP}} \end{bmatrix} := \operatorname{argmin}_{b, \beta} \left\{ \|y - Zb - X\beta\|_2^2 + b^T (\lambda_Q Q + \lambda_R I) b \right\}$$

This problem is “equivalent” to the LMM formulation

- 1 $y = Zb + X\beta + \varepsilon$, where β is a vector of fixed effects and b a vector of random effects,
- 2 $\varepsilon \sim \mathcal{N}(0, \sigma^2 I)$,
- 3 $b \sim \mathcal{N}\left(0, (\sigma_Q^2 Q + \sigma_R^2 I)^{-1}\right)$,
- 4 $\lambda_Q, \lambda_R, \sigma, \sigma_Q$ and σ_R are connected via
 $\lambda_Q = \sigma^2 / \sigma_Q^2, \lambda_R = \sigma^2 / \sigma_R^2.$

Simulation scheme

SIMULATED SIGNAL




Laplacian : Q_{true}

„Invertible Laplacian” :
 $\tilde{Q}_{true} := Q_{true} + 0.001 \cdot I$

True signal used in simulation:

$$b^{true} \sim N(0, \sigma_b^2 \tilde{Q}_{true}^{-1})$$

ESTIMATION



Laplacian of distorted graph was used to find the estimate,




$\hat{b}$

MSEr defined as

$$\text{MSEr} := E \left[\frac{\|\hat{b} - b_{true}\|_2^2}{\|b_{true}\|_2^2} \right]$$

as a measure of estimation accuracy

Simulation scheme – distorted connectivity matrices



Simulation scheme

Three methods compared:

- 1 ridge: $\lambda_Q := 0$ (connectivity information is not used)
- 2 naive: $\lambda_R := 0$, $Q \rightarrow \tilde{Q}$ (only λ_Q is selected)
- 3 riPEER (both lambdas are selected in an adaptive way)


Axis of the plot

- 1 X axis:


$$\text{diss}(A^{\text{true}}, A^{\text{obs}}) := \frac{\text{number of removed/added connections}}{\text{number of all nonzero connections in } A^{\text{true}}}$$

- 2 Y axis: $MSEr := \mathbb{E} \left[\frac{\|\hat{b} - b^{\text{true}}\|_2^2}{\|b^{\text{true}}\|_2^2} \right]$.


Simulation results



Simulation results




Simulation results



Results: HIV study

- ① Association between cortical thickness and speed of information processing
- ② 66 considered brain's regions
- ③ $N = 199$ individuals

Results: Speed of Information Processing



- Cortical regions: lingual[L], precentral[L], superiorparietal[L], lateralorbitofrontal[R], precentral[R], superiorparietal[R], supramarginal[R], medialorbitofrontal[L], posteriorcingulate[R]


Non-Gaussian distributions

- $y_i \sim$ member of an Exponential family of distribution
- Consider an optimization problem of the form

$$\operatorname{argmin}_{b, \beta} \left\{ \underbrace{-2 \loglik(y; \beta, b)}_{\text{model fit term}} + \underbrace{g_{\lambda}(b)}_{\text{penalty on } b} \right\}.$$

- $g_{\lambda}(b) := \lambda_Q b^T Q b + \lambda_R \|b\|_2^2$
- λ_Q and λ_R are selected based on the equivalence with GLMM

HIV data: HIV+ vs. HIV-



Contributions: riPEER and griPEER

- 1 **Brain connectivity informed regularization methods for regression**, *Statistics in Biosciences*, April 2019, Volume 11, Issue 1, 47–90
mdpeer R package is available online at
<https://CRAN.R-project.org/package=mdpeer>
- 2 **Connectivity-Informed Adaptive Regularization Under non-Gaussian Design**, doi:
<https://doi.org/10.1101/322420>
griPEER Matlab software is available online at
<https://github.com/dbrzyski/griPEER>

Conclusions and Discussion

- ➊ Regularization methods using a priori information in brain imaging setting in a principled way.
- ➋ Additional (external) information quantified via a Laplacian matrix.
- ➌ Penalty parameters chosen in an adaptive way via ML/REML.
- ➍ The more knowledge one has about informative structure, the more specific one can be in defining the penalty matrix.
- ➎ Use of structural connectivity to inform the associations between cortical thickness and either NP outcomes (continuous) or HIV status (binary).

Funding: NIMH R01MH108467

Collaborators



Damian Brzyski
Technical Univ. of
Wroclaw, Poland



Marta Karas
Johns Hopkins
University



Timothy Randolph
Fred Hutchinson
C. R. Center



Mario Dziedzic
Indiana University



Joaquin Goni
Purdue
University

- *Brain Connectivity-Informed Regularization Methods for Regression, Statistics in Biosciences (2017),*
<https://doi.org/10.1007/s12561-017-9208-x>
- **mdpeer R package** is available online at
<https://CRAN.R-project.org/package=mdpeer>



Beau Ances
Washington University

References

- ❶ M. Karas, D. Brzyski, M. Dziedzic, J. Goni, D. A. Kareken, T. W. Randolph, J. Harezlak, *Brain connectivity-informed regularization methods for regression*, Stat Biosci (2019). <https://doi.org/10.1007/s12561-017-9208-x>.
- ❷ T. W. Randolph, J. Harezlak, Z. Feng, *Structured penalties for functional linear models – partially empirical eigenvectors for regression*, Electronic Journal of Statistics **6** (2012), 323–353.
- ❸ C. Li, H. Li, *Network-constrained Regularization and Variable Selection for Analysis of Genomic Data*, JBioinformatics **24** (2008), no. 9, 1175–1182.
- ❹ F. Chung, *Laplacians and the Cheeger Inequality for Directed Graphs*, Annals of Combinatorics **9** (2005), no. 1, 1–19.
- ❺ E. Demidenko, *Mixed Models: Theory and Applications*, Wiley **9** (2004).

Pictures sources:

- ❶ FreeSurfer, <http://www.opensourceimaging.org/project/freesurfer/>
- ❷ Athinoula A. Martinos Center, <http://www.martinos.org/neurorecovery/technology.htm>
- ❸ PsyPost, <http://www.psypost.org/2014/04/atypical-brain-connectivity-associated-with-autism-spectrum-disorder-24472>